

Tailor's Bunion (Bunionette)





American College of
Foot and Ankle Surgeons®

What Is a Tailor's Bunion?

Tailor's bunion, also called a bunionette, is a prominence of the fifth metatarsal bone at the base of the little toe. The metatarsals are the five long bones of the foot. The prominence that characterizes a tailor's bunion occurs at the metatarsal "head," located at the far end of the bone where it meets the toe. Tailor's bunions are not as common as bunions, which occur on the inside of the foot, but they are similar in symptoms and causes.

Why is it called a tailor's bunion? The deformity received its name centuries ago, when tailors sat cross-legged all day with the outside edge of their feet rubbing on the ground. This constant rubbing led to a painful bump at the base of the little toe.



Causes

Often a tailor's bunion is caused by an inherited faulty mechanical structure of the foot. In these cases, changes occur in the foot's bony framework, resulting in the development of an enlargement. The fifth metatarsal bone starts to protrude outward, while the little toe moves inward. This shift creates a bump on the outside of the foot that becomes irritated whenever a shoe presses against it.

Sometimes a tailor's bunion is actually a bony spur (an outgrowth of bone) on the side of the fifth metatarsal head.

Regardless of the cause, the symptoms of a tailor's bunion are usually aggravated by wearing shoes that are too narrow in the toe, producing constant rubbing and pressure.

Symptoms

The symptoms of tailor's bunions include redness, swelling, and pain at the site of the enlargement. These symptoms occur when wearing shoes that rub against the enlargement, irritating the soft tissues underneath the skin and producing inflammation.

Diagnosis

Tailor's bunion is easily diagnosed because the protrusion is visually apparent. X-rays may be ordered to help the foot and ankle surgeon determine the cause and extent of the deformity.

Tailor's Bunion

(Bunionette) *continued*American College of
Foot and Ankle Surgeons®**Non-Surgical Treatment**

Treatment for tailor's bunion typically begins with non-surgical therapies. Your foot and ankle surgeon may select one or more of the following:

- **Shoe modifications.** Choose shoes that have a wide toe box, and avoid those with pointed toes or high heels.
- **Padding.** Bunionette pads placed over the area may help reduce pain.
- **Oral medications.** Nonsteroidal anti-inflammatory drugs (NSAIDs), such as ibuprofen, may help relieve the pain and inflammation.
- **Icing.** An ice pack may be applied to reduce pain and inflammation. Wrap the pack in a thin towel rather than placing ice directly on your skin.

- **Injection therapy.** Injections of corticosteroid may be used to treat the inflamed tissue around the joint.

- **Orthotic devices.** In some cases, custom orthotic devices may be provided by the foot and ankle surgeon.

When Is Surgery Needed?

Surgery is often considered when pain continues despite the above approaches. In selecting the procedure or combination of procedures for your case, the foot and ankle surgeon will take into consideration the extent of your deformity based on the x-ray findings, your age, your activity level, and other factors. The length of the recovery period will vary, depending on the procedure or procedures performed. ■

Kenneth R. Meisler, D.P.M., P.L.L.C. & Associates
210 E. 86th Street Suite 402
New York, NY 10028
(212) 628-4444
<http://nypodiatry.net>

American College of
Foot and Ankle Surgeons®

This information has been prepared by the Consumer Education Committee of the American College of Foot and Ankle Surgeons, a professional society of over 6,000 foot and ankle surgeons. Members of the College are Doctors of Podiatric Medicine who have received additional training through surgical residency programs.

The mission of the College is to promote superior care of foot and ankle surgical patients through education, research and the promotion of the highest professional standards.

Copyright © 2010, American College of Foot and Ankle Surgeons • www.FootHealthFacts.org